

| RESEARCH ARTICLE

Article DOI: 10.21474/JNAVES01/150

DOI URL: <http://dx.doi.org/10.21474/JNAVES01/150>

**PESTICIDE RESIDUES IN AGRICULTURAL SOILS: ENVIRONMENTAL RISKS, SOIL
HEALTH, AND SUSTAINABLE MANAGEMENT STRATEGIES**

Sana M. Khan, Arjun P. Verma and Olivia R. Bennett

Corresponding Author: Sana M. Khan

| ABSTRACT

Pesticides play an important role in modern agriculture by helping farmers control insects, weeds, fungi, and other organisms that damage crops. However, inappropriate application, excessive use, and inadequate disposal can contribute to pesticide residues in agricultural soils. Depending on their chemical properties and environmental conditions, these substances may persist in soil, move into groundwater, affect non-target organisms, and influence important biological processes. This paper examines the sources, behavior, environmental consequences, and management of pesticide residues in agricultural soils. It uses a narrative literature review approach to synthesize established research and guidance from recognized scientific and regulatory organizations. The review discusses the influence of soil texture, organic matter, moisture, temperature, and microbial activity on pesticide persistence and movement. It also explores the effects of residues on soil microorganisms, beneficial insects, water quality, and agricultural biodiversity. Sustainable management options include integrated pest management, appropriate pesticide selection, application according to label instructions, crop rotation, vegetative buffer zones, and regular environmental monitoring. The paper emphasizes that pesticide behavior varies considerably among compounds and locations, so risk assessment must be based on reliable evidence rather than general assumptions. It concludes that protecting agricultural soils requires cooperation among farmers, researchers, agricultural extension services, and regulatory authorities.

| KEYWORDS

Pesticide residues, agricultural soil, soil health, environmental pollution, integrated pest management, biodiversity, sustainable agriculture, groundwater protection.

| ARTICLE INFORMATION

RECEIVED: 04 July 2026

ACCEPTED: 10 August 2026

PUBLISHED: September 2026

Introduction:-

Agriculture is essential for food production, rural employment, and economic development. Farmers face many challenges in protecting crops from insects, weeds, plant diseases, and other pests. Pesticides are commonly used to reduce crop losses and support agricultural productivity. When selected and applied appropriately, they can be valuable tools in crop protection. Nevertheless, pesticides may also create environmental concerns when they are used incorrectly or when their movement and persistence are not adequately understood. Some pesticides or their breakdown products can remain in soil after application. Others may degrade relatively quickly, depending on their chemical structure and environmental conditions. Agricultural soil is a living system containing minerals, organic matter, microorganisms, insects, earthworms, plant roots, and many other organisms. These components support nutrient cycling, water regulation, plant growth, and ecosystem stability. Chemical contamination may affect some of these functions, although the magnitude of the effect depends on the pesticide, its concentration, exposure duration, and local conditions.

Pesticide residues may reach soil directly during spraying or indirectly through contaminated irrigation water, pesticide drift, or the use of certain contaminated materials. Rainfall and irrigation can influence whether residues remain near the soil surface, move deeper into the soil, or enter nearby water bodies. Understanding pesticide behavior is therefore important for maintaining soil health and protecting agricultural ecosystems. This paper reviews the environmental risks associated with pesticide residues and identifies practical management approaches that can support sustainable crop production.

Copyright: © 2026 the Author(s). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) 4.0 license (<https://creativecommons.org/licenses/by/4.0/>). Published by Jana Publication and Research LLP.

Objectives of the Study:-

The main objectives of this paper are:-

1. To explain the major sources of pesticide residues in agricultural soils.
2. To examine the environmental factors that influence pesticide persistence and movement.
3. To discuss potential effects on soil organisms, biodiversity, and water quality.
4. To describe methods used to monitor pesticide residues.
5. To evaluate sustainable pesticide-management practices.
6. To identify research priorities for reducing pesticide-related environmental risks.

Review of Literature:-

Research on pesticide behavior demonstrates that pesticides differ considerably in their persistence, mobility, toxicity, and interactions with soil components. Some compounds bind strongly to soil organic matter, while others are more mobile in water. Environmental conditions such as temperature, moisture, soil acidity, and microbial activity can influence degradation.

The Food and Agriculture Organization of the United Nations and the World Health Organization have developed guidance on pesticide management and the evaluation of pesticide risks. Their work emphasizes responsible use, proper handling, and the importance of considering effects on human health and the environment. The United States Environmental Protection Agency also evaluates pesticides according to their potential risks and establishes requirements for approved uses. Such regulatory approaches demonstrate the importance of assessing individual substances rather than assuming that all pesticides behave in the same way. Scientific literature also highlights integrated pest management as an approach that combines monitoring, prevention, biological control, cultural practices, and carefully selected chemical treatments when necessary. Integrated pest management can help reduce unnecessary pesticide applications while maintaining effective crop protection. Despite progress in pesticide regulation and management, environmental monitoring remains important because agricultural conditions vary across regions and because pesticide mixtures and degradation products can complicate risk assessment.

Research Methodology:-

Research Design:-

This paper adopts a narrative literature review design. It synthesizes established scientific knowledge concerning pesticide residues, soil processes, environmental risks, and sustainable agricultural management.

Sources of Information:-

The review framework draws on publications and guidance from recognized organizations, including the Food and Agriculture Organization of the United Nations, the World Health Organization, the United States Environmental Protection Agency, and peer-reviewed scientific literature on soil contamination and pesticide behavior.

Thematic Analysis:-

The material is organized into five themes: pesticide sources, environmental persistence, effects on soil ecosystems, monitoring approaches, and sustainable management practices.

Limitations:-

This paper does not report original laboratory testing, field sampling, or experimental data. It is a narrative review rather than a systematic review or meta-analysis. Its conclusions therefore provide a general synthesis and should not be interpreted as findings from a new field investigation.

Sources of Pesticide Residues in Agricultural Soils:-

Direct Application to Crops:-

Pesticides may reach soil when spray droplets fall beyond the target crop, when chemicals are applied directly to soil, or when treated plant material decomposes. Some pesticides are specifically designed for soil application, while others reach soil unintentionally during crop spraying. The quantity entering soil depends on application technique, weather conditions, crop structure, and the chemical properties of the product.

Excessive or Incorrect Application:-

Applying pesticides more frequently or at higher rates than permitted can increase environmental exposure. Incorrect equipment calibration, unsuitable spraying conditions, and failure to follow label instructions may also contribute to off-target movement. Training and proper application equipment can help farmers reduce unnecessary losses and improve the effectiveness of pest-control measures.

Contaminated Irrigation Water:-

Water used for irrigation may contain pesticide residues originating from agricultural runoff or other sources. Repeated irrigation with contaminated water may introduce residues into soil over time. Water-quality monitoring is particularly relevant in agricultural areas where surface water receives drainage from treated fields.

Improper Storage and Disposal:-

Leaking containers, spills, and inappropriate disposal of pesticide products can create localized contamination. Washing application equipment near wells, streams, or drainage channels may also introduce chemicals into the environment. Secure storage, spill-prevention measures, and disposal through approved local systems are important elements of responsible pesticide management.

Atmospheric Drift and Deposition:-

Pesticide droplets or particles may move beyond the intended application area through wind. Depending on the product and conditions, some substances may also move through the atmosphere and later settle onto soil or vegetation. Suitable weather conditions, appropriate equipment, and compliance with product instructions can help reduce off-target movement.

Factors Influencing Pesticide Persistence in Soil:-**Soil Organic Matter:-**

Organic matter can influence pesticide adsorption and availability. Some compounds bind strongly to organic matter, reducing their immediate mobility, although binding does not necessarily eliminate long-term environmental concerns. Soils with different organic matter levels may therefore behave differently when exposed to the same pesticide.

Soil Texture and Mineral Composition:-

Clay minerals and other soil components can affect how chemicals bind to soil particles. Sandy soils may allow greater movement of some mobile compounds under certain conditions, whereas finer-textured soils may retain some substances more strongly. However, the behavior of a particular pesticide cannot be predicted from texture alone. Chemical properties and local conditions must also be considered.

Temperature and Moisture:-

Temperature and soil moisture influence microbial activity and chemical reactions. Under suitable conditions, microorganisms can break down certain pesticides. Extremely dry, waterlogged, hot, or cold conditions may change the rate of degradation in different ways. The effect varies according to the compound and the organisms present in the soil.

Soil Acidity and Alkalinity:-

Soil pH can influence pesticide stability, solubility, adsorption, and degradation. Some pesticides break down more rapidly under acidic or alkaline conditions, while others behave differently. Understanding the relationship between soil chemistry and pesticide properties can improve environmental risk assessment.

Chemical Structure:-

Pesticides differ in their molecular structures and chemical characteristics. These differences influence how long they remain in the environment, whether they bind to soil, and how readily they move through water.

For this reason, the environmental persistence of one pesticide should not be generalized to an entire pesticide group without supporting evidence.

Effects on Soil Health and Biodiversity:-**Soil Microorganisms:-**

Soil bacteria and fungi participate in decomposition, nutrient cycling, and the maintenance of soil structure. Some pesticide exposures may affect the activity or composition of these microbial communities. The extent of the effect depends on the chemical, concentration, exposure period, soil characteristics, and the organisms present. Certain changes may be temporary, while other situations may require further investigation. Maintaining soil organic matter, reducing unnecessary chemical exposure, and monitoring soil biological indicators can support soil health.

Earthworms and Other Soil Invertebrates:-

Earthworms help mix organic matter into soil and contribute to soil structure and water movement. Other invertebrates perform important ecological functions, including decomposition and supporting food webs. Some pesticides can adversely affect non-target invertebrates at particular exposure levels. Risk assessment should therefore consider organisms beyond the intended pest species.

Nutrient Cycling:-

Microorganisms help transform nutrients into forms that plants can use. If a pesticide affects organisms involved in these processes, nutrient cycling may also be influenced. However, effects are not identical across compounds or soil conditions. Reliable assessment requires appropriate measurements and a clear understanding of the chemical exposure involved.

Pollinators and Beneficial Organisms:-

Pesticide residues in soil may affect some organisms that spend part of their life cycle underground or come into contact with contaminated soil. Other beneficial organisms may be exposed through spray drift, contaminated plants, or water. Reducing unnecessary applications and choosing targeted control methods can help limit exposure of beneficial species.

Agricultural Biodiversity:-

Healthy agricultural ecosystems contain a variety of organisms that contribute to pollination, pest regulation, decomposition, and soil fertility. Chemical exposure may alter some of these relationships. Biodiversity-friendly farming practices, including habitat protection and integrated pest management, can complement responsible pesticide use.

Movement of Pesticides into Water Resources:-**Surface Runoff:-**

Rainfall or irrigation can carry pesticide residues from agricultural land into nearby streams, ponds, and other surface-water bodies. The risk may increase when water flows rapidly across treated fields. Vegetated buffer strips, soil conservation, appropriate application timing, and runoff management can help reduce the movement of some pesticides into surface water.

Leaching into Groundwater:-

Certain pesticides can move downward through the soil with infiltrating water. Whether this occurs depends on chemical mobility, soil properties, rainfall, irrigation, and groundwater depth. Groundwater contamination is particularly important because groundwater may supply drinking water and irrigation. Monitoring should focus on areas where the characteristics of the pesticide and the local environment indicate elevated risk.

Effects on Aquatic Organisms:-

Pesticides entering water may affect aquatic organisms, including invertebrates, fish, and other wildlife. The effects depend on the substance, concentration, exposure duration, and sensitivity of the species. Preventing contamination at its source is generally preferable to relying only on treatment after pesticides have entered water bodies.

Monitoring and Detection of Pesticide Residues:-**Soil Sampling:-**

Soil sampling provides information about the presence and distribution of pesticide residues. Sampling plans should account for differences in land use, soil type, application history, and depth. Samples should be collected, stored, transported, and documented using appropriate procedures to maintain data quality.

Laboratory Analysis:-

Laboratories use analytical techniques to identify and quantify pesticide residues. Methods may include gas chromatography and liquid chromatography coupled with suitable detectors or mass spectrometry. The appropriate method depends on the pesticide being investigated, the expected concentration, and the characteristics of the soil sample.

Quality Assurance:-

Reliable results require validated analytical methods, suitable reference materials, calibration, blanks, and other quality-control procedures. Detection limits and measurement uncertainty should be reported where relevant. Without proper quality assurance, residue measurements may be difficult to compare or interpret accurately.

Environmental Risk Assessment:-

Finding a pesticide residue does not automatically mean that harmful effects are occurring. Risk assessment considers the chemical's toxicity, concentration, exposure route, duration, environmental fate, and the organisms that may be exposed. Results should be compared with relevant scientific benchmarks and applicable regulations, where available.

Long-Term Monitoring:-

Repeated sampling can help identify trends and determine whether management changes are associated with lower environmental exposure. Long-term programs are especially valuable where pesticides are used frequently or where water resources and sensitive habitats may be affected.

Sustainable Strategies for Pesticide Management:-

Integrated Pest Management:-

Integrated pest management (IPM) combines multiple methods to prevent and manage crop pests. These may include crop rotation, resistant varieties, field monitoring, biological control, habitat management, and targeted chemical treatments when justified. IPM does not necessarily eliminate pesticide use. Instead, it aims to ensure that treatments are selected according to pest identification, monitoring results, economic thresholds where appropriate, and environmental considerations.

Responsible Pesticide Selection:-

Farmers should select products approved for the intended crop and pest and follow all label instructions. Product choice should consider effectiveness, environmental fate, potential non-target effects, and applicable restrictions. Pesticides should not be selected solely on the basis of price or immediate effectiveness.

Proper Application Techniques:-

Correct equipment calibration, appropriate nozzles, suitable weather conditions, and avoidance of unnecessary spraying can reduce off-target exposure. Farmers should follow required protective measures and buffer distances, observe re-entry and pre-harvest intervals, and comply with local laws.

Crop Rotation and Cultural Practices:-

Crop rotation can disrupt the life cycles of certain pests and reduce dependence on repeated use of the same control method. Timely cultivation, appropriate planting schedules, sanitation, and resistant crop varieties may also help manage pests. The effectiveness of these methods depends on the crop, pest species, local climate, and farm system.

Biological Control:-

Biological control uses natural enemies or other biological approaches to manage pests. Depending on the system, this may involve beneficial insects, microorganisms, or conservation of natural predators. Biological control requires correct identification and appropriate management. Introducing non-native organisms without adequate assessment can create new environmental problems.

Vegetative Buffer Zones:-

Vegetated strips along field boundaries and waterways may help reduce runoff, soil erosion, and movement of some pesticide residues. Their effectiveness depends on vegetation, width, slope, soil conditions, and rainfall. Buffer zones should be designed according to local guidance and should not be treated as a complete substitute for responsible pesticide application.

Role of Farmers, Researchers, and Policymakers:-

Farmer Education:-

Farmers need accessible information about pesticide labels, application equipment, storage, spill prevention, and environmental risks. Training should use practical demonstrations and be available in locally understood languages. Agricultural extension services can help farmers select suitable alternatives and interpret local pest-monitoring information.

Scientific Research:-

Researchers can investigate pesticide persistence, interactions between pesticide mixtures, effects on non-target organisms, and the effectiveness of alternative pest-management approaches. Studies should include different soil types, climatic conditions, and agricultural systems to avoid conclusions based on a narrow range of environments.

Regulatory Oversight:-

Regulatory authorities play an important role in evaluating pesticide safety, approving uses, establishing requirements, and monitoring compliance. Regulations should be supported by reliable scientific evidence and updated as new information becomes available. Effective enforcement also requires appropriate laboratory capacity, trained personnel, and communication with farming communities.

Community Participation:-

Local communities can contribute to identifying environmental concerns, protecting water resources, and supporting responsible agricultural practices. Cooperation among farmers, researchers, and local authorities can improve awareness and encourage practical solutions.

Challenges and Limitations:-

Several challenges affect the prevention and management of pesticide residues in agricultural soils:-

Financial constraints: Soil testing, improved equipment, and alternative pest-management systems may involve costs that small farms cannot easily afford.

Limited technical knowledge: Farmers may not have access to reliable guidance on pesticide properties, application techniques, and environmental protection.

Complex environmental behavior: Pesticide persistence and movement vary among compounds and locations, making general predictions difficult.

Limited monitoring capacity: Some regions lack sufficient laboratories, trained staff, or regular sampling programs.

Pest-management pressures: Crop losses and the need for reliable harvests may encourage repeated pesticide use where alternative methods are unavailable.

Regulatory differences: Requirements and approved products vary across jurisdictions, so farmers need guidance that reflects local law.

Addressing these challenges requires practical education, affordable monitoring, suitable alternatives, and stronger cooperation between agricultural and environmental institutions.

Future Research Directions:-

Future research should focus on developing practical and evidence-based methods to reduce pesticide-related environmental risks.

Important priorities include:-

1. Investigating the persistence of commonly used pesticides under different soil and climatic conditions.
 2. Improving methods for detecting pesticide residues and relevant degradation products.
 3. Examining the combined effects of pesticide mixtures on soil organisms.
 4. Evaluating integrated pest management under different crop-production systems.
 5. Developing affordable residue-monitoring approaches for agricultural communities.
 6. Assessing the effectiveness of buffer strips and other measures for reducing runoff.
 7. Investigating relationships between pesticide exposure, soil biological indicators, and long-term soil health.
 8. Improving farmer access to information about approved products and safe handling.
 9. Comparing the environmental and economic outcomes of alternative pest-control strategies.
 10. Strengthening communication between laboratories, agricultural extension services, and policymakers.
- Research should report methods, uncertainties, and limitations transparently to support informed decision-making.

Recommendations:-

Based on the review, the following recommendations are proposed:-

- Promote integrated pest management to reduce unnecessary pesticide applications.
- Encourage farmers to follow approved product labels and local regulations.
- Improve the safe storage, transport, handling, and disposal of pesticides.
- Strengthen soil and water monitoring in areas where contamination risks are elevated.
- Protect waterways through appropriate buffer zones and runoff management.
- Support farmer training in pesticide safety and environmental protection.
- Encourage crop rotation, resistant varieties, and suitable biological control methods.
- Expand access to reliable soil-testing laboratories and technical advice.
- Improve coordination among agricultural agencies, environmental authorities, and research institutions.
- Evaluate pesticide-management interventions according to local evidence and practical farm conditions.

Conclusion:-

Pesticides remain important tools for protecting crops, but their use must be managed carefully to prevent avoidable harm to agricultural soils and surrounding ecosystems. Pesticide residues may influence soil organisms, nutrient cycling, biodiversity, and water quality, although the level of risk depends on the individual chemical, exposure conditions, and environmental setting. Sustainable management requires more than simply reducing the quantity of chemicals applied. It involves selecting appropriate products, using them correctly, monitoring environmental conditions, adopting integrated pest management, and preventing contamination during storage and disposal. Farmers, researchers, agricultural extension services, and regulatory authorities all have roles in improving pesticide management. Greater access to education, affordable monitoring, and practical alternatives can help farming communities protect soil health while maintaining productive agriculture. In conclusion, responsible pesticide use and evidence-based environmental management are essential for sustainable agricultural development. Protecting soil today can support productive farming, ecological stability, and food security for future generations.

Author Contributions:-

The actual authors should specify their individual contributions to the literature review, analysis, drafting, and revision. The sample author names above are fictional and do not represent verified contributors.

Conflict of Interest:-

The actual authors should disclose any financial or personal relationships that could influence the work.

Funding Statement:-

Funding information should be added only after confirmation from the actual authors. If the study received no external funding, the authors may state this according to the journal's requirements.

Data Availability Statement:-

No original dataset was generated for this narrative literature review.

Ethical Considerations:-

This review did not involve original experiments on animals or human participants. Any future field sampling or laboratory investigation should follow relevant institutional, environmental, and regulatory requirements.

References:-

1. Food and Agriculture Organization of the United Nations (FAO), & World Health Organization (WHO). (2014). International Code of Conduct on Pesticide Management. FAO. <https://www.fao.org/pest-and-pesticide-management/en/>
2. Food and Agriculture Organization of the United Nations (FAO). (n.d.). Integrated Pest Management. FAO. <https://www.fao.org/pest-and-pesticide-management/ipm/integrated-pest-management/en/>
3. United States Environmental Protection Agency (EPA). (n.d.). Pesticide environmental stewardship. <https://www.epa.gov/pesticide-stewardship>
4. World Health Organization (WHO). (n.d.). Chemical safety: Pesticides. <https://www.who.int/health-topics/chemical-safety>
5. Food and Agriculture Organization of the United Nations (FAO). (n.d.). Plant Production and Protection. <https://www.fao.org/plant-production-protection/en/>